


SEARCH  
ADVANCED SEARCH

DONATE HELP CONTACT AHA SIGN IN HOME

Feedback Subscriptions Archives Search Table of Contents

**Hypertension** **citeTrack** Let the article find you!



Hypertension, Vol 19, 249-254, Copyright © 1992 by American Heart Association

## ARTICLES

### Characterization of a monoclonal antibody specific for human active renin

WM Zuo, RE Pratt, CH Heusser, JP Bews, MM de Gasparo and VJ Dzau

Division of Cardiovascular Medicine, Stanford University School of Medicine, Calif.

We have identified and characterized an anti-human renin monoclonal antibody R1-20-5 that is selective for human active renin. R1-20-5 binds active renin with a dissociation constant ( $K_d$ ) of  $2.5 \times 10^{-7}$  M/l and inhibits renin enzymatic activity with an inhibitory constant ( $IC_{50}$ ) of  $1.4 \times 10^{-8}$  M/l. R1-20-5 competes with a synthetic renin inhibitor for binding with renin, demonstrating further that it is binding to or close to the active site. This antibody does not bind prorenin in human plasma or recombinant prorenin expressed by L-929 fibroblasts transfected with human renin gene. Furthermore, trypsin activation of prorenin resulted in immunoreactivity of the activated prorenin toward the antibody. In addition, an immunoaffinity column of R1-20-5 coupled to Sepharose retained active renin but had a low affinity for prorenin. A sensitive and rapid solid phase radioimmunoassay for active renin was developed using a "sandwich" technique employing R1-20-5 and a second non-active site-directed monoclonal antibody to human renin. Renin levels in human plasma samples were determined by the standard enzymatic assay, and by the direct radioimmunoassay for active renin, before and after trypsin activation. Trypsin treatment of plasma resulted in parallel increases in both the plasma renin enzymatic activity and in the plasma active renin concentration as measured by the direct radioimmunoassay. Overall, plasma immunoreactive active renin concentration correlated significantly with plasma renin enzymatic activity ( $r = 0.96$ ,  $p$  less than 0.001). In summary, the monoclonal antibody R1-20-5 is selective for human active renin and should be a very useful tool for studies of the active enzyme in humans.

#### This Article

- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)

#### Services

- ▶ [Email this article to a friend](#)
- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Download to citation manager](#)
- ▶ [Cited by other online articles](#)
- ▶ [Request Permissions](#)

#### PubMed

- ▶ [PubMed Citation](#)
- ▶ [Articles by Zuo, W. M.](#)
- ▶ [Articles by Dzau, V. J.](#)

### This article has been cited by other articles:



**Am. J. Physiol: Heart and Circulatory Physiology**


▶ HOME

J. J. Saris, F. H. M. Derkx, R. J. A. De Bruin, D. H. W. Dekkers, J. M. J. Lamers, P. R. Saxena, M. A. D. H. Schalekamp, and A. H. Jan Danser

**High-affinity prorenin binding to cardiac man-6-P/IGF-II receptors precedes proteolytic activation to renin**

Am J Physiol Heart Circ Physiol, April 1, 2001; 280(4): H1706 - 1715.

[Abstract] [Full Text]



## Circulation

[HOME](#)

A. H. J. Danser, C. A. M. van Kesteren, W. A. Bax, M. Tavenier, F. H. M. Derkx, P. R. Saxena, and M. A. D. H. Schalekamp

### **Prorenin, Renin, Angiotensinogen, and Angiotensin-Converting Enzyme in Normal and Failing Human Hearts : Evidence for Renin Binding**

Circulation, July 1, 1997; 96(1): 220 - 226.

[\[Abstract\]](#) [\[Full Text\]](#)

[HOME](#) [HELP](#) [FEEDBACK](#) [SUBSCRIPTIONS](#) [ARCHIVE](#) [SEARCH](#) [TABLE OF CONTENTS](#)

[HYPERTENSION](#)

[ART, THRO, VASC BIO](#)



[ALL AHA JOURNALS](#)

[CIRCULATION](#)

[CIRCULATION RESEARCH](#)

[STROKE](#)

**Copyright © 1992 by the American Heart Association.**



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for   

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract  Sort 

Text Version

All: 1 Review: 0 

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

☐ 1: Hypertension. 1992 Mar;19(3):249-54.[Related Articles, Links](#)

## Characterization of a monoclonal antibody specific for human active renin.

**Zuo WM, Pratt RE, Heusser CH, Bews JP, de Gasparo MM, Dzau VJ.**

Division of Cardiovascular Medicine, Stanford University School of Medicine, Calif.

We have identified and characterized an anti-human renin monoclonal antibody R1-20-5 that is selective for human active renin. R1-20-5 binds active renin with a dissociation constant ( $K_d$ ) of  $2.5 \times 10^{-7}$  M/l and inhibits renin enzymatic activity with an inhibitory constant ( $IC_{50}$ ) of  $1.4 \times 10^{-8}$  M/l. R1-20-5 competes with a synthetic renin inhibitor for binding with renin, demonstrating further that it is binding to or close to the active site. This antibody does not bind prorenin in human plasma or recombinant prorenin expressed by L-929 fibroblasts transfected with human renin gene. Furthermore, trypsin activation of prorenin resulted in immunoreactivity of the activated prorenin toward the antibody. In addition, an immunoaffinity column of R1-20-5 coupled to Sepharose retained active renin but had a low affinity for prorenin. A sensitive and rapid solid phase radioimmunoassay for active renin was developed using a "sandwich" technique employing R1-20-5 and a second non-active site-directed monoclonal antibody to human renin. Renin levels in human plasma samples were determined by the standard enzymatic assay, and by the direct radioimmunoassay for active renin, before and after trypsin activation. Trypsin treatment of plasma resulted in parallel increases in both the plasma renin enzymatic activity and in the plasma active renin concentration as measured by the direct radioimmunoassay. Overall, plasma immunoreactive active renin concentration correlated significantly with plasma renin enzymatic activity ( $r = 0.96$ ,  $p$  less than 0.001). In summary, the monoclonal antibody R1-20-5 is selective for human active renin and should be a very useful tool for studies of the active enzyme in humans.

PMID: 1548051 [PubMed - indexed for MEDLINE]

Display Abstract  Sort [Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)

Department of Health &amp; Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)